

学校编码：10384

学号：7463

廈門大學

博士学位论文

海岸带地下水循环与海底地下水排泄研究

Study on Groundwater Cycle and Submarine
Groundwater Discharge in the Coastal Zone

刘花台

指导教师：高爱国

专业名称：海洋地质

答辩日期：2014年9月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1. 经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

摘 要

海岸带地下水循环和海底地下水排泄是全球水循环的重要组成部分，近年来成为陆-海相互作用研究的主要内容之一。海底地下水排泄不单是一个水循环过程，还是一个地球化学循环过程，其输送大量营养盐、碳、金属等物质到海洋中，在海洋物质收支平衡中具有重要意义。本论文的主要目标是了解福建沿海的地下水循环特点和地下水入海通量，为此，在福建省科技计划项目和国家自然科学基金的资助下，选择厦门五缘湾、闽江河口和漳州前湖湾为研究区域，通过现场观测、调查和取样，并采用同位素示踪技术、电法探测和水文地球化学等研究方法和手段，对水体滞留时间、海底地下水排泄量、河水与海水混合速率、地下水与河水补排关系、地下水、河水及海水的混合比例、不同类型含水层中地下水循环特点、海底泉成因、海岸带地下水循环模式等问题开展了研究。

(1) 厦门五缘湾

五缘湾是厦门岛内的一个小海湾，其海岸属于砂质类型海岸，海岸带主体由未固结的晚更新世海积-冲积物和全新世海积物组成。五缘湾的水体滞留时间和地下水入海通量是重点研究内容。

借助 ^{224}Ra 和 ^{226}Ra 半衰期的差异，利用 ^{224}Ra 和 ^{226}Ra 比值计算得到五缘湾水体的表观年龄在0.6~2.4 d之间，平均滞留时间为1.4 d，其中湾顶水体滞留时间较长。该结果与用纳潮量方法计算的滞留时间1.2 d比较接近。

地下水和海水中 ^{224}Ra 与 ^{226}Ra 活度的比值特征表明，输入五缘湾的地下水，70.2%源自冲积层中的孔隙潜水，29.8%源自海积层中的孔隙潜水。通过构建 ^{224}Ra 和 ^{226}Ra 的质量平衡模型（箱式模型），计算获得地下水输入的 ^{224}Ra 和 ^{226}Ra 通量分别为 $5.17 \times 10^6 \text{ Bq/d}$ 和 $5.28 \times 10^6 \text{ Bq/d}$ ，转化为海底地下水排泄量为 $4.16 \times 10^5 \text{ m}^3/\text{d}$ 和 $4.73 \times 10^5 \text{ m}^3/\text{d}$ 。

(2) 闽江河口区

闽江河口区是一个基岩海岸和砂质海岸共存的海岸类型，河水、地下水和海水三者相互作用明显。通过枯水期和丰水期的调查和取样，重点对河水与地下水的补排关系、河水与海水的混合速率、河水、地下水及海水的混合比例，以及地下水入海通

量进行了研究。

河流悬浮颗粒上 ^{224}Ra 和 ^{226}Ra 的解吸程度随盐度升高而增大，在盐度 $17.0\sim 18.0$ 时， ^{224}Ra 和 ^{226}Ra 基本上达到完全解吸。当河流悬浮颗粒上的 ^{224}Ra 和 ^{226}Ra 完全解吸以后，混合水中的 ^{224}Ra 和 ^{226}Ra 活度随着离岸距离的增加而呈指数衰减，由此计算获得涨潮时段内河水与海水的混合速率为 $140.2\sim 142.5\text{ m/h}$ 。

水样的 $\delta^{2}\text{H}$ 、 $\delta^{18}\text{O}$ 散点图较好地反映出地下水和河水的起源与演化，以及地下水与河水的相互补排关系。在枯水期，闽江南岸地下水接受当地降水补给后又经历了一定的蒸发作用，而闽江北岸地下水接受当地降水补给后基本上没有受到后期蒸发作用的影响；在丰水期，闽江两岸地下水均接受了大气降水的充分补给，蒸发作用轻微。无论枯水期还是丰水期，闽江河水都是接受两岸地下水的补给；在枯水期，闽江南岸地下水经历一定的蒸发作用后再补给河水，而在丰水期两岸地下水基本上接受降水补给后又快速补给河水。

以河水、地下水和海水为端元构建三端元混合模型，计算获得丰水期闽江河口区河水、地下水和海水的混合比率分别是 34.2% 、 10.3% 和 55.5% 。以闽江的月平均入海径流量为依据，计算获得丰水期和枯水期闽江河口区地下水的入海通量分别是 $58.1\times 10^6\text{ m}^3/\text{d}$ 和 $7.5\times 10^6\text{ m}^3/\text{d}$ ，前者约是后者的 7.7 倍。进一步计算获得丰水期地下水输入闽江河口的溶解无机氮、活性磷酸盐、溶解性硅酸盐通量分别是 $12.4\times 10^6\text{ mol/d}$ 、 $0.5\times 10^6\text{ mol/d}$ 和 $10.8\times 10^6\text{ mol/d}$ 。

(3) 漳州前湖湾

前湖湾是比较典型的海岸沙丘发育的砂质海岸，岸线较为平直，海滩平坦宽阔，非常有利于研究海岸带的地下水循环。而且，前湖湾在潮间带发育海底泉，这在国内是很少见的，泉水的来源至今还是个谜。为此，通过调查和取样对海岸带地下水循环特点和模式、海底泉的形成开展了研究。

近岸海积层-风积层互层构成的含水层系统，主要接受降水的垂直补给，含水层渗透性和富水性均很好，水循环速度较快，直接向海域排泄。水化学成分(Cl/Br 比、 Ca/Mg 比、 Ca/SO_4 比)表明，该含水系统孔隙水中仍部分保留海侵时期的海水组分特征。表层风积物中的孔隙水，受现代海水的影响，表现出部分海水的特征。受人为过量开采地下水的影响，近岸局部地段出现不同程度的海水入侵现象。滨海平

原冲洪积层中的孔隙水，径流比较缓慢，溶滤作用和蒸发浓缩作用同时存在。风化残积层中的裂隙-孔隙水，其化学组成变化很大，水循环速度明显受局部地形的影响和控制。新近纪玄武岩-砂岩组裂隙-孔隙水，阳离子 Na^+ 、 Mg^{2+} 和 Ca^{2+} 毫克当量百分数都超过25%，阴离子则以 HCO_3^- 为主，说明其水循环速度远高于冲洪积层孔隙水和风化残积层裂隙-孔隙水，这与其所处地势较高，以及玄武岩垂直裂隙发育、砂岩呈半固结状等有直接关系。

高密度电法揭示海底泉在海滩上的出露直径约为5.5~6.0 m，泉眼处垂向上的延伸深度约是2.5~3.0 m；海底泉的补给路径不与海岸线垂直，而是偏向其西北方向，该方向与基底燕山期侵入岩中断裂带的发育方向比较吻合。

海底泉与新近纪玄武岩-砂岩组裂隙-孔隙水化学组成的相似性、海底泉异常高的 ^{222}Rn 活度等证据表明，海底泉的补给区可能是位于莲花山一带的新近纪玄武岩-砂岩含水岩组，该含水岩组的裂隙-孔隙水通过下伏燕山期花岗岩的断裂带输送给海底泉。

潮间带孔隙水盐度分布很不均一，明显受到海底泉和陆源淡水排泄的影响。垂直岸线的潮间带孔隙水盐度分布特征表明，潮汐泵海水小循环、盐水楔海水大循环和陆源地下淡水排泄共同构成了前海湾近岸基本的水循环模式，海底泉排泄使得该水循环模式变得复杂化。

潮间带浅表层（70 cm以浅）孔隙水的主要离子和营养盐组成表明，潮间带孔隙水循环速度很快，其地球化学过程比较简单，化学反应微弱，主要是地下淡水与海水的物理混合作用。

综上所述，福建省海岸带类型比较复杂，不同类型海岸带的地下水循环特点和海底地下水排泄量存在差异。氢氧稳定同位素示踪法、镭和氡放射性同位素示踪法、电阻率法、水文地球化学方法各有其优势，针对海岸带类型特点和具体地质、构造及水文地质条件，适宜采用不同方法，或者综合应用多种方法对海岸带的地下水循环和海底地下水排泄进行研究。

关键词：地下水循环；海底地下水排泄；镭同位素；氢氧稳定同位素；电阻率法

Abstract

Coastal groundwater cycle and submarine groundwater discharge is an important part of global water cycle, it recently has been become one of the research focus of land-ocean interactions in the coastal zone. Submarine groundwater discharge is not only a water cycle process but also a geochemical cycle process, these processes deliver massive dissolved material such as nutrients, carbon and metal into the ocean, which plays a significant role in the budget of marine chemical material. The aim of this thesis is to discuss the groundwater cycle characteristics and groundwater discharge into ocean, funded by the province scientific plan of Fujian and the national natural science foundation of China.

This thesis presents the study on the related contents of groundwater cycle and submarine groundwater discharge in the Wuyuan bay of Xiamen, Minjiang estuary and Qianhu bay of Zhangzhou, such as water mass residence time, groundwater discharge flux, recharge relationship of riverine water and groundwater, mixing ratio between groundwater, riverine water and seawater, groundwater cycle characteristics in different aquifers, cause of formation for submarine spring, groundwater cycle model in the coastal zone, which are discussed by the approach of field investigation, sampling, isotope tracer, electrical resistivity detecting and hydrogeochemical method.

(1) Wuyuan bay of Xiamen

Wuyuan bay is a small bay located in Xiamen island, its coast belongs to sandy coast, composed by unconsolidated Pleistocene's marine-alluvial deposit and Holocene's marine deposit. This part focuses on the seawater residence time and groundwater discharge flux in Wuyuan bay.

Based on the difference of ^{224}Ra and ^{226}Ra half life, apparent water age in Wuyuan bay was calculated to be from 0.6~2.4 d, with average residence time 1.4 d and longer residence time in the bay head. This average residence time is

close to 1.2 d calculated by tidal prism approach.

The feature of ^{224}Ra and ^{226}Ra ratio value in groundwater and seawater indicates that 70.2% of groundwater into Wuyuan bay is attributed to the pore water from alluvium, another 20.8% is attributed to the pore water from marine deposit. Through establishing ^{224}Ra and ^{226}Ra mass balance model(box model), ^{224}Ra and ^{226}Ra fluxes driven by groundwater were calculated to be $5.17 \times 10^6 \text{ Bq/d}$ and $5.28 \times 10^6 \text{ Bq/d}$, respectively. Transformed from these ^{224}Ra and ^{226}Ra fluxes, submarine groundwater discharge fluxes are $4.16 \times 10^5 \text{ m}^3/\text{d}$ and $4.73 \times 10^5 \text{ m}^3/\text{d}$, respectively.

(2) Minjiang estuary

The interaction between riverine water, groundwater and seawater in the Minjiang estuary, where rocky coast and sandy coast coexist, is very clear. Based on field investigation and sampling in dry season and wet season, respectively, relationship between riverine water and groundwater, mixing rate of riverine water and groundwater, mixing ratio between riverine water, groundwater and seawater, and groundwater discharge flux into ocean are key discussion contents of this part.

^{224}Ra and ^{226}Ra desorbing amount from suspending riverine particles increases with increasing salinity, ^{224}Ra and ^{226}Ra basically reach the complete desorption at the salinity of 17.0~18.0. After ^{224}Ra and ^{226}Ra desorbed completely from suspending riverine particles, ^{224}Ra and ^{226}Ra activities in the mixing water exponentially decay with increasing offshore distance, based on this rule, mixing rates between riverine water and seawater in rising tide period were calculated to be 140.2~142.5 m/h.

$\delta^2\text{H}$ and $\delta^{18}\text{O}$ scatter diagram from water samples better revealed the origin and evolution of groundwater and riverine water, and recharge relationship between groundwater and riverine water. In dry season, groundwater at the south bank of Min jiang river experiences somewhat evaporation after receiving the recharge of

rainfall, while groundwater at the north bank of Minjiang river seldom suffer from the influence of evaporation. In wet season, the groundwater at both banks of Minjiang river receive the recharge of rainfall, with slight evaporation. No matter whether dry season or wet season, Minjiang river's water receive the recharge from both banks of Minjiang river; in dry season, the groundwater at both banks of Minjiang river recharge riverine water after experiencing somewhat evaporation, but in wet season, the groundwater at both banks of Minjiang river rapidly recharge riverine water as soon as it receives the recharge of rainfall.

In wet season, through establishing three end-member mixing model, the mixing ratio between riverine water, groundwater and seawater was calculated to be 34.2%, 10.3% and 55.5%, respectively. Then, based on the monthly average runoff of Minjiang river, groundwater discharge fluxes into ocean were calculated to be 58.1×10^6 m³/d for the wet season and 7.5×10^6 m³/d for the dry season, respectively. The former is 7.7 times larger than the latter. According to the groundwater fluxes, the dissolved inorganic nitrogen, reactive phosphate and soluble silicate to the ocean in wet season were estimated to be 12.4×10^6 mol/d, 0.5×10^6 mol/d and 10.8×10^6 mol/d, respectively.

(3) Qianhu bay of Zhangzhou

Qianhu bay is a typical sandy coast, with better developmental sand dune, straighter shoreline, and flat and wide beach, this kind of coast is very suitable to discussing the groundwater cycle. Moreover, submarine spring emerges at the beach of Qianhu bay, which is a rather rare natural phenomenon in China's coast zone, its origin is a mystery by far. In this part, the characteristics and pattern of groundwater cycle, origin of submarine spring were discussed through investigation and sampling.

Inshore aquifer system constituted by marine deposit and eolian deposit, with better permeability and water-abundance, and rapid water cycle velocity, mainly receives the vertical recharge of rainfall, and its groundwater directly flow into the

ocean. Hydrochemical component (Cl/Br, Ca/Mg, Ca/SO₄) indicate that the pore water of this aquifer system still partly shows the feature of seawater formed during transgression. The pore water in surface eolian deposit partly presents the feature of seawater influenced by modern seawater. Partial seawater intrusion derived from overpumping occurred at different levels. The pore water from alluvium of coastal plain has obvious leaching and concentration with slow runoff. Fracture-pore water in elurium has different water cycle velocity controlled by local landform, its chemical component varies largely. For the fracture-pore water of Neogene's basalt-sand rock formation, the percentage of milliequivalent of cation Na+Mg²⁺ and Ca²⁺ exceed 25% with dominated anion HCO₃⁻ show that its water cycle velocity is higher than above pore water of alluvium and fracture-pore water of elurium, which directly related to the elevated landform, developmental vertical fissure in basalt and unconsolidated sand rock.

Electrical resistivity approach revealed that the submarine spring diameter of 5.5~6.0 m emerge at the beach, and vertical extending depth is about 2.5~3.0 m at the mouth of spring; the recharge pathway of submarine spring is not perpendicular to the shoreline, but it tends to northwest direction, this direction is consistent with the developmental direction of fault zone formed in Yanshan epoch's intrusive rock.

Some evidence such as the similarity of chemical constituent between submarine spring and fracture-pore water from the Neogene's basalt-sand rock formation, unusually high ²²²Rn activity of spring water indicate that the recharge area of submarine spring probably comes from the Neogene's basalt-sand rock formation located in Lianhua mountain, and the fracture-pore water of this water-bearing formation recharge the spring through the fault zone of underlying Yanshan epoch's intrusive rock.

The salinity distribution of pore water in the intertidal zone is not very uniform influenced by submarine spring and fresh terrestrial groundwater discharge. The

characteristics of salinity distribution of pore water perpendicular to shoreline indicate that the smaller tidal pump cycle, bigger salt water wedge cycle and fresh terrestrial groundwater discharge together constitute the fundamental water cycle pattern of Qianhu bay, and submarine spring makes this water cycle pattern complicated.

The component of main ions and nutrients of surface pore water(<70 cm) in the intertidal zone show that pore water cycles rapidly, and its geochemical process is simple, chemical reaction is weak, major process is the mechanical mix of fresh groundwater and seawater.

In conclusion, the coastal type of Fujian is more various, different type of coast has different groundwater cycle and submarine groundwater discharge. The different research methods such as tracer of stable hydrogen and oxygen isotopes, tracer of radioactive Ra and Rn, electrical resistivity, hydrogeochemical method, have their own advantages and disadvantages. For the different coastal type and specific geology, tectonic and hydrogeologic condition, single method or multi-method should be suitable to apply in the study on groundwater cycle and submarine groundwater discharge in the coastal zone.

Keywords: groundwater cycle; submarine groundwater discharge; Radium isotope; stable hydrogen and oxygen isotopes; electrical resistivity method

参考资料

- Andersen M S, Jakobsen V N R, Postma D. Geochemical processes and solute transport at the seawater/freshwater interface of a sandy aquifer[J]. *Geochimica et Cosmochimica Acta*, 2005, 69(16): 3979-3994.
- Back W, Hanshaw B B, Pylar T E, et al. Geochemical significance of groundwater discharge in Caleta Xel Ha, Quintana Roo, Mexico[J]. *Water Resources Research*, 1979, 15: 1521-1535.
- Basu A R, Jacobsen S B, Poreda R J, et al. Large groundwater strontium flux to the oceans from the Bengal basin and the marine strontium isotope record[J]. *Science*, 2001, 293: 1470-1473.
- Beck A J, Rapaglia J P, Cochran J K. Radium mass-balance in Jamaica Bay, NY: Evidence for a substantial flux of submarine groundwater[J]. *Marine Chemistry*, 2007a, 106: 419-441.
- Beck A J, Tsukamoto Y, Tovar-Sanchez A. Importance of geochemical transformations in determining submarine groundwater discharge-derived trace metal and nutrient fluxes[J]. *Applied Geochemistry*, 2007b, 22: 477-490.
- Bokuniewicz H J. Groundwater seepage into Great South Bay, New York[J]. *Estuarine and Coastal Marine Science*, 1980, 10: 504-508.
- Bone S E, Charette M A, Lamborg C H, et al. Has submarine groundwater discharge been overlooked as a source of mercury to coastal waters?[J] *Environmental Science & Technology*, 2007, 41: 3090-3095.
- Bouchaou L, Michelot J L, Vengosh A, et al. Application of multiple isotopic and geochemical tracers for investigation of recharge, salinization, and residence time of water in the Souss – Massa aquifer, southwest of Morocco[J]. *Journal of Hydrology*, 2008, 352: 267-287.
- Bratton J F, Bölke J K, Krantz D E, et al. Flow and geochemistry of groundwater beneath a back-barrier lagoon: The subterranean estuary at Chincoteague Bay, Maryland, USA[J]. *Marine Chemistry*, 2009, 113: 78-92.
- Buddemeier R.W. Groundwater flux to the ocean: definitions, data, applications, uncertainties [C].Buddemeier R.W. (ed), *Groundwater discharge in the coastal zone: Proceedings of an international symposium, LOICZ IGBP*. Texel, the Netherlands: LOICZ, 1996, 16-21.
- Burnett W C. Offshore springs and seeps are focus of working group[J]. *EOS, American Geophysical Union*, 1999, 80: 13-15.
- Burnett W C, Taniguchi M, Oberdorfer J. Measurement and significance of the direct discharge of groundwater into the coastal zone[J]. *Journal of Sea Research*, 2001, 46: 109-116.
- Burnett W C, Bokuniewicz H, Huettel M, et al. Groundwater and pore water inputs to the coastal zone[J]. *Biogeochemistry*, 2003, 66: 3-33.
- Burnett W C, Aggarwal P K, Aureli A, et al. Quantifying submarine groundwater discharge in the coastal zone via multiple methods [J]. *Science of the Total Environment*, 2006, 367: 498-543.
- Cai W J, Wang Y C, Krest J, et al. The geochemistry of dissolved inorganic carbon in a surficial groundwater aquifer in North Inlet, South Carolina, and the carbon fluxes to the coastal ocean[J]. *Geochimica et Cosmochimica Acta*, 2003, 67: 631-639.
- Charette M A, Buesseler K O, Andrews J E. Utility of radium isotopes for evaluating the input and transport of groundwater-derived nitrogen to a Cape Cod estuary[J]. *Limnology and Oceanography*, 2001, 46(2): 465-470.
- Charette M A, Sholkovitz E R. Oxidative precipitation of groundwater-derived ferrous iron in the subterranean estuary of a coastal bay[J]. *Geophysical Research Letters*, 2002, 29, doi: 10.1029/2001gl014512.
- Charette M A, Buesseler K O. Submarine groundwater discharge of nutrients and copper to an urban subestuary of Chesapeake Bay (Elizabeth River)[J]. *Limnology and Oceanography*, 2004, 49: 376-85.
- Charette M A, Sholkovitz E R, Hansell C M. Trace element cycling in a subterranean estuary: Part 1. Geochemistry of the permeable sediments[J]. *Geochimica et Cosmochimica Acta*, 2005, 69: 2095-2109.
- Charette M A, Shokovitz E R. Trace element cycling in a subterranean eastury: Part 2. Geochemistry of the pore water[J]. *Geochimica et Cosmochimica Acta*, 2006, 70: 811-826.

Craig H. Isotopic variations in meteoric waters[J]. *Science*, 1961, 133(3465): 1702-1703.

Crotwell A M, Moore W S. Nutrient and radium fluxes from submarine groundwater discharge to Port Royal Sound, South Carolina[J]. *Aquatic Geochemistry*, 2003, 9: 191-208.

Dai M, Yin Z, Meng F, et al. Spatial distribution of riverine DOC inputs to the ocean[J]. *Current Opinion in Environmental Sustainability*, 2012, 4: 170-178.

Dansgaard W. Stable isotopes in precipitation[J]. *Tellus*, 1964, 16(4): 436-468.

Dettmann E H. Effect of water residence time on annual export and denitrification of nitrogen in estuaries: a model analysis[J]. *Estuaries*, 2001, 24: 481-490.

Dronkers J, Zimmerman J T F. Some principles of mixing in tidal lagoons. p. 107-117. *Proceedings of the International Symposium on Coastal Lagoons*. Bordeaux, France, 1982.

Dulaiova H, Burnett W C. Evaluation of the flushing rates of Apalachicola Bay, Florida via natural geochemical tracers[J]. *Marine Chemistry*, 2008, 109: 395-408.

Duncan T, Shaw T J. The mobility of rare earth elements and redox sensitive elements in the ground-water/seawater mixing zone of a shallow coastal aquifer[J]. *Aquatic Geochemistry*, 2003, 9: 233-255.

Dzhamalov R G, Safronova T I. On estimating chemical discharge into the world ocean with groundwater[J]. *Water Resources*, 2002, 29(6): 626-631.

Einsiedl F. Sea-water/groundwater interactions along a small catchment of the European Atlantic coast[J]. *Applied Geochemistry*, 2012, 27: 73-80.

Ferrarin C, Rapaglia J, Zaggia L. Coincident application of a mass balance of radium and a hydrodynamic model for the seasonal quantification of groundwater flux into the Venice Lagoon, Italy[J]. *Marine Chemistry*, 2008, 112: 179-188.

Garcia-Solsona E, Masque P, Garcia-Orellana J, et al. Estimating submarine groundwater discharge around Isola La Cura, northern Venice Lagoon (Italy), by using the radium quartet[J]. *Marine Chemistry*, 2008, 109: 292-306.

Garrison G H, Glenn C R. Measurement of submarine groundwater discharge in Kahana Bay, Oahu, Hawaii [J]. *Limnology and Oceanography*, 2003, 48(2): 920-928.

Gonneea M E, Morris P J. New perspectives on radium behavior within a subterranean estuary[J]. *Marine Chemistry*, 2008, 109: 250-267.

Goni M A, Gardner L R. Seasonal dynamics in dissolved organic carbon concentrations in a coastal water-table aquifer at the forest-marsh interface[J]. *Aquatic Geochemistry*, 2003, 9:209-232.

Gu H Q, Moore W S, Zhang L, et al. Using radium isotopes to estimate the residence time and the contribution of submarine groundwater discharge (SGD) in the Changjiang effluent plume, East China Sea[J]. *Continental Shelf Research*, 2012, 35: 95-107.

Hanshaw B B, Back W. A geochemical hypothesis for dolomitization by groundwater[J]. *Economic Geology*, 1971, 66: 710-724.

Hougham A L, Moran S B. Water mass ages of coastal ponds estimated using ^{223}Ra and ^{224}Ra as tracers[J]. *Marine Chemistry*, 2007, 105: 194-207.

Johannesson K H, Burdige D J. Balancing the global oceanic neodymium budget: Evaluating the role of groundwater[J]. *Earth and Planetary Science Letters*, 2007, 253: 129-42.

Johannes R E. The ecological significance of the submarine discharge of ground water[J]. *Marine Ecology Progress Series*, 1980, 3: 365-73.

Kazemi G A. Editor's Message: Submarine groundwater discharge studies and the absence of hydrogeologists[J]. *Hydrogeology Journal*, 2008, 16: 201-204.

Kim G, Ryu J W, Hwang D W. Radium tracing of submarine groundwater discharge (SGD) and associated nutrient fluxes in a highly-permeable coastal zone, Korea[J]. *Marine Chemistry*, 2008, 109: 307-317.

Kohout F A. Submarine springs: A neglected phenomenon of coastal hydrology[J]. *Hydrology*, 1966, 26: 391-413.

Kontar E A, Ozorovich Y R. Geo-electromagnetic survey of the fresh/salt water interface in the coastal southeastern Sicily[J]. *Continental Shelf Research*, 2006, 26: 843-851.

Ladouche B, Weng P. Hydrochemical assessment of the Rochefort marsh: Role of surface and groundwater in the hydrological functioning of the wetland[J]. *Journal of Hydrology*, 2005, 314: 22-42.

Lambert M J, Burnett W C. Submarine groundwater discharge estimates at a Florida coastal site based on continuous radon measurements[J]. *Biogeochemistry*, 2003, 66: 55-73.

Lee D R. A device for measuring seepage flux in lakes and estuaries [J]. *Limnology and Oceanography*, 1977, 22: 140-147.

Lee Jin-yong, Cho Byung-wook. Submarine groundwater discharge into the coast revealed by water chemistry of man-made undersea liquefied petroleum gas cavern[J]. *Journal of Hydrology*, 2008, 36: 195-206.

Lin I T, Wang C H, You C F, et al. Deep submarine groundwater discharge indicated by tracers of oxygen, strontium isotopes and barium content in the Pingtung coastal zone, southern Taiwan[J]. *Marine Chemistry*, 2010, 122: 51-58.

Lin In-tian, WANG Chung-ho, LIN Saulwood, et al. Groundwater – seawater interactions off the coast of southern Taiwan: Evidence from environmental isotopes[J]. *Journal of Asian Earth Sciences*, 2011, 41: 250-262.

Liu Q, Dai M, Chen W, et al. How significant is submarine groundwater discharge and its associated dissolved inorganic carbon in a river-dominated shelf system[J]. *Biogeosciences*, 2012, 9: 1777-1795.

Martin J H, Fitzwater S. Iron deficiency limits phytoplankton growth in the northeast Pacific subarctic[J]. *Nature*, 1988, 331: 341-343.

Martin J B, Hartl K M, Corbett D R, et al. A multi-level pore-water sampler for permeable sediments[J]. *Journal of Sedimentary Research*, 2003, 73(1): 128-132.

Milliman J D. Production and accumulation of calcium carbonate in the ocean: Budget of a nonsteady state[J]. *Global Biogeochemistry Cycles*, 1993, 7: 927-57.

Men W, Wang Fenfen, Zhang Yusheng, et al. Determining coastal mixing rates of western Taiwan Strait using ²²⁴Ra[J]. *Journal of Radioanalytical and Nuclear Chemistry*, 2012, DOI 10.1007/s10967-012-1853-4.

Monsen N E, Cloern J E, Luzzas L V. A comment on the use of flushing time, residence time, and age as transport time scales[J]. *Limnology and Oceanography*, 2002, 47(5): 1545-1553.

Moore W S. Sampling radium-228 in the deep ocean[J]. *Deep-Sea Research*, 1976, 23: 647-651.

Moore W S. Large groundwater inputs to coastal waters revealed by ²²⁶Ra enrichments[J]. *Nature*, 1996, 380: 612-614.

Moore W S. The effects of groundwater input at the mouth of the Ganges-Brahmaputra Rivers on barium and radium fluxes to the Bay of Bengal[J]. *Earth and Planetary Science Letters*, 1997, 150:141-150.

Moore W S. The subterranean estuary: a reaction zone of groundwater and sea water[J]. *Marine Chemistry*, 1999, 65: 111-125.

Moore W S. Determining coastal mixing rates using radium isotopes[J]. *Continental Shelf Research*, 2000a, 20: 1993-2007.

Moore W S. Ages of continental shelf waters determined from Ra-223 and Ra-224[J]. *Journal of Geophysical Research*, 2000b, 105: 22117-22.

Moore W S. Sources and fluxes of submarine groundwater discharge delineated by radium isotopes[J]. *Biogeochemistry*, 2003, 66: 75-93.

Moore W S, Blanton J O, Joye S B. Estimates of flushing times, submarine groundwater discharge, and nutrient fluxes to Okatee Estuary, South Carolina[J]. *Journal of Geophysical Research*, 2006, 111, doi: 10.1029/2005jc003041.

Moore W S. Fifteen years experience in measuring ²²⁴Ra and ²²³Ra by delayed-coincidence counting[J]. *Marine Chemistry*, 2008, 109: 188-197.

Moore W S. The Effect of Submarine Groundwater Discharge on the Ocean[J]. *The Annual Review of Marine Science*, 2010, 2: 59-88.

Oberdorfer J A. Hydrogeologic modeling of submarine groundwater discharge: comparison to other quantitative methods[J]. *Biogeochemistry*, 2003, 66: 159-169.

Paulsen R J, Smith C F, O'Rourke D, et al. Development and evaluation of an ultrasonic ground water seepage

meter[J]. *Ground Water*, 2001, 39: 904-911.

Peterson R N, Burnett W C, Taniguchi M, et al. Radon and radium isotope assessment of submarine groundwater discharge in the Yellow River delta, China[J]. *Journal of Geophysical Research: Oceans* (1978-2012), 2008b, 113: C09021.

Peterson R N, Burnett W C, Taniguchi M, et al. Determination of Transport Rates in the Yellow River – Bohai Sea Mixing Zone via Natural Geochemical Tracers[J]. *Continental Shelf Research*, 2008a, 28: 2700-2707.

Povinec P P, Bokuniewicz H, Burnett W C. et al. Isotope tracing of submarine groundwater discharge offshore Ubatuba, Brazil: results of the IAEA – UNESCO SGD project [J]. *Journal of Environmental Radioactivity*, 2008, 99: 1596-1610.

Povinec P P, Aggarwal P K, Aureli A, et al. Characterisation of submarine groundwater discharge offshore south-eastern Sicily. *Journal of Environmental Radioactivity*[J], 2006, 89: 81-101.

Rapaglia J, Ferrarin C, Zaggia L. Investigation of residence time and groundwater flux in Venice Lagoon: comparing radium isotope and hydrodynamical models[J]. *Journal of Environmental Radioactivity*, 2010, 101: 571-581.

Robinson C, Li L, Barry D A. Effect of tidal forcing on subterranean estuary[J]. *Advances in Water Resources*, 30: 851-865.

Roxburgh I S. Thermal infrared detection of submarine springs associated with the Plymouth Limestone[J]. *Hydrology Science Journal*, 1985, 30: 185-96.

Sanders C J, Santos I R, Barcellos R, et al. Elevated concentrations of dissolved Ba, Fe and Mn in a mangrove subterranean estuary[J]. *Continental Shelf Research*, 2012, 43: 86-94.

Sanford L, Boicourt W, Rives S. Model for estimating tidal flushing of small embayments[J], *Journal of Waterway Port Coastal and Ocean Engineering*, 1992, 118: 635-654.

Santos I R, Burnett W C, Chanton J, et al. Nutrient biogeochemistry in a Gulf of Mexico subterranean estuary and groundwater-derived fluxes to the coastal ocean[J]. *Limnology and Oceanography*, 2008, 53 (2): 705-718.

Santos I R, Burnett W C, Dittmar T, et al. Tidal pumping drives nutrient and dissolved organic matter dynamics in a Gulf of Mexico subterranean estuary[J]. *Geochimica et Cosmochimica Acta*, 2009, 73: 1325-1339.

Santos I R, Eyre B D, Huettel M. The driving forces of porewater and groundwater flow in permeable coastal sediments: A review[J]. *Estuarine, Coastal and Shelf Science*, 2012, 98: 1-15.

Schiavo M A, Hauser S, Povinec P P. Stable isotopes of water as a tool to study groundwater – seawater interactions in coastal south-eastern Sicily[J]. *Journal of Hydrology*, 2009, 364: 40-49.

Schmidt A, Santos I R, Burnett W C, et al. Groundwater sources in a permeable coastal barrier: Evidence from stable isotopes[J]. *Journal of Hydrology*, 2011, 406: 66-72.

Shaw T J, Moore W , Kloepper J, et al. The flux of barium to the coastal waters of the southeast-ern USA: The importance of submarine groundwater discharge[J]. *Geochimica et Cosmochimica Acta*, 1998, 62: 3047-3054.

Sholkovitz E, Herbold C, Charette M. An automated dye-dilution based seepage meter for the time-series measurement of submarine groundwater discharge[J]. *Limnology and Oceanography*, 2003, 1: 16-28.

Smiths L, Zawadzki W. A hydrogeologic model of submarine groundwater discharge: Florida intercomparison experiment [J]. *Biogeochemistry*, 2003, 66: 95-110.

Spiteri C, Slomp C P, Charette M A, et al. Flow and nutrient dynamics in a subterranean estuary (Waquoit Bay, MA, USA): field data and reactive transport modeling[J]. *Geochimica et Cosmochimica Acta*, 2008, 72: 3398-2412.

Stieglitz T, Rapaglia J, Bokuniewicz H. Estimation of submarine groundwater discharge from bulk ground electrical conductivity measurements[J]. *Journal Geophysics Research*, 2008, 113, doi: 10.1029/2007jc004499.

Su N, Du J Z, Moore W S, et al. An examination of groundwater discharge and the associated nutrient fluxes into the estuaries of eastern Hainan Island, China using ²²⁶Ra[J]. *Science of the Total Environment*, 2011, 409: 3909-3918.

Sun Y, Torgersen T. The effects of water content and Mn-fiber surface conditions on ²²⁴Ra measurement by ²²⁰Rn emanation[J]. *Marine Chemistry*, 1998, 62: 299-306.

Taniguchi M, Burnett W C, Cable J E, et al. Investigation of submarine groundwater discharge [J]. *Hydrological Processes*, 2002(16): 2115-2129.

Taniguchi M, Burnett W C, et al. Spatial and temporal distributions of submarine groundwater discharge rates obtained from various types of seepage meters at a site in the Northeastern Gulf of Mexico[J]. *Biogeochemistry*, 2003, 66: 35-53.

Taniguchi M, Ishitobi T, Shimada J. Dynamics of submarine ground-water discharge and freshwater-seawater interface[J]. *Journal of Geophysical Research*, 2006, 111: C01008. doi:10.1029/2005JC002924.

Tse K C, Jiao J J. Estimation of submarine groundwater discharge in Plover Cove, Tolo Harbour, Hong Kong by ^{222}Rn [J]. *Marine Chemistry*, 2008, 111(3-4): 160-170.

Ueda S, Suzumura M, Sumi E. Denitrification in a seashore sandy deposit influenced by groundwater discharge[J]. *Biogeochemistry*, 2003, 63: 187-205.

Werner A D, Bakker M, Post V E et al. Seawater intrusion processes, investigation and management: Recent advances and future challenges[J]. *Advances in Water Resources*, 2013, 51: 3-26.

Wilson A M. Fresh and saline groundwater discharge to the ocean: A regional perspective[J]. *Water Resources Research*, 2005, 41, doi: 10.1029/2004wr003399.

Windom H, Niencheski F. Biogeochemical processes in a freshwater-seawater mixing zone in permeable sediments along the coast of Southern Brazil[J]. *Marine Chemistry*, 2003, 81: 121-130.

Windom H L, Moore W S, Niencheski L F H et al. Submarine groundwater discharge: A large, previously unrecognized source of dissolved iron to the South Atlantic Ocean[J]. *Marine Chemistry*, 2006, 102: 252-266.

Younger P L. Submarine groundwater discharge[J]. *Nature*, 1996, 382: 121-122.

Zektser I S, Ivanov V A, Meskheteli A V. The problem of direct groundwater discharge to the seas[J]. *Journal of Hydrology*, 1973, 20: 1-36.

Zektser I S, Everett L G, Dzhamalov R G. *Submarine Groundwater*[C]. Boca Raton, FL: CRC Press, 2007.

Zhang J, Mandal A K. Linkages between submarine groundwater systems and the environment [J]. *Current Opinion in Environmental Sustainability*, 2012, (4): 219-226.

Zimmerman J T F. Estuarine residence times, p.75-84. In B. Kjerfve [ed.] *Hydrodynamics of estuaries*[M]. CRC Press, 1988.

陈中笑, 程军, 郭品文等. 中国降水稳定同位素的分布特点及其影响因素[J]. *大气科学学报*, 2010, 33(6): 667-679.

戴志忠. 福州地区水文特性分析[J]. *水利科技*, 2004, 3: 9-11.

谷现平, 聂新恕, 周江等. 利用高密度电法仪探测地下水流速流向[J]. *中国煤炭地质*, 2010, 22(增刊): 83-85.

国家环境保护总局编. *水和废水监测分析方法*[M]. 北京: 中国环境科学出版社, 2002.

郭占荣, 黄磊, 刘花台等. 镭同位素示踪隆教湾的海底地下水排泄[J]. *地球学报*, 2008, 29(5): 647-652.

郭占荣, 黄磊, 袁晓婕等. 用镭同位素评价九龙江河口区的地下水输入[J]. *水科学进展*, 2011, 22(1): 118-125.

郭占荣, 李开培, 袁晓婕等. 用氡-222评价五缘湾的地下水输入[J]. *水科学进展*, 2012, 23(2): 263-270.

郭占荣, 马志勇, 章斌等. 采用 ^{222}Rn 示踪胶州湾的海底地下水排泄及营养盐输入[J]. *地球科学—中国地质大学学报*, 2013, 38(5): 1073-1080.

黄卿团, 付萍, 胡哲明等. 福建闽江断裂带第四纪活动特征[J]. *震灾防御技术*, 2007, 2(4): 346-353.

黄一民, 章新平, 唐方雨等. 长沙大气降水中稳定同位素变化及过量氡指示水汽来源[J]. *自然资源学报*, 2013, 28(11): 1945-1954.

黄奕普, 谢永臻, 陈性保等. 海水中 ^{224}Ra 快速富集和测定的新方法[J]. *厦门大学学报*, 2001, 40(3): 699-705.

季仲强, 胡丹, 翁焕新等. 近岸海域 ^{226}Ra 的时空变化与海底地下水排泄(SGD)估算[J]. *地球化学*, 2012, 41(1): 15-22.

贾国东, 黄国伦. 海底地下水排泄: 重要的海岸带陆海相互作用过程[J]. *地学前缘*, 2005, 12: 29-35.

李福林, 陈学群, 张奇等. 莱州湾东岸海底地下水的水文地球化学特征及其地球化学演化模式[J]. *海洋*

科学进展, 2005, 23(2): 190-196.

李开培. 五缘湾的地下水输入研究[D]. 厦门大学硕士学位论文, 2011.

李海龙, 万力, 焦赳赳. 海岸带水文地质学研究中的几个热点问题[J]. 2011, 26(7): 685-694.

李建成. 福建沿海地区含水岩组水文地质条件分析[J]. 泉州师范学院学报(自然科学), 2002, 20(6): 54-57.

李艳平. 镭同位素的地球化学行为及其海洋学意义[D]. 厦门大学硕士学位论文, 2004.

廖小青, 刘贯群, 袁瑞强等. 黄河农场地区地下水入海量FEFLOW软件数值模拟[J]. 海洋科学进展, 2005, 23(4): 446-451.

林晨. 福州盆地工程地质特征及其勘察与施工[J]. 矿产保护与利用, 2002, 2(5): 52-54.

刘贯群, 叶玉玲, 袁瑞强等. 近年胶州湾陆源SGD及其营养盐输送[J]. 海洋环境科学, 2007, 26(6): 510-513.

刘广山. 海洋放射性核素测量方法[M]. 北京: 海洋出版社, 2006.

刘花台, 郭占荣, 袁晓婕等. 用镭同位素评价海水滞留时间及海底地下水排泄[J]. 地球科学—中国地质大学学报, 2013, 38(3): 599-606.

刘花台, 郭占荣, 高爱国等. 闽江河口区水体中镭的分布特征及河水与海水的混合速率[J]. 吉林大学学报(地球科学版), 2013, 43(6): 1966-1971.

刘花台, 郭占荣. 海底地下水排泄的研究进展[J]. 地球科学进展, 2014, 29(7): 33-44.

刘茜. 海底地下水排泄及其对近海无机碳循环的影响[D]. 厦门大学博士学位论文, 2012.

刘延锋, 江贵荣, 靳孟贵等. 新疆焉耆盆地水环境氢氧同位素特征及其指示作用[J]. 地质科技情报, 2009, 28(6): 89-93.

刘维坤. 厦门岛及其邻近地区新构造运动基本特征[J]. 台湾海峡, 1983, 2(1): 82-90.

刘忠方, 田立德, 姚檀栋等. 中国大气降水中 ^{18}O 的空间分布[J]. 科学通报, 2009, 54(6): 804-811.

门武. 镭同位素示踪的黄海和东海海洋学研究[D]. 厦门大学博士论文, 2008.

门武, 刘广山, 陈敏等. 中国东海的 ^{224}Ra [J]. 地球科学-中国地质大学学报, 2011, 36(6): 999-1008.

邱汉学, 郑西来, 张效龙等. 黄河农场地区地下水入海通量的数值分析[J]. 海洋地质动态, 2003, 19(3): 28-33.

邱晓晖. Mn-纤维富集和测定天然水中Ra-226 方法的研究[D]. 厦门大学硕士学位论文, 1988.

苏妮. 镭同位素示踪的近岸水体混合和海底地下水排泄[D]. 华东师范大学博士学位论文, 2013.

许博超. 天然镭同位素富集和测定方法及对河口混合过程的示踪研究[D]. 中国海洋大学博士论文, 2011.

宋献方, 夏军, 于静洁等. 应用环境同位素技术研究华北典型流域水循环机理的展望[J]. 地理科学进展, 2002, 21(6): 527-537.

宋献方, 刘相超, 夏军等. 基于环境同位素技术的怀沙河流域地表水和地下水转化关系研究[J]. 中国科学 D辑: 地球科学, 2007, 37(1): 102-110.

田玉昆, 刘怀山, 张晶等. 高密度电法寻找地下水在即墨地区的应用[J]. 工程地球物理学报, 2008, 5(6): 670-674.

厦门地质工程勘察院. 厦门市区域水文地质调查报告(1:10万)[R]. 1999.

许苏清, 潘伟然, 张国荣等. 浔江湾海水交换时间的计算[J]. 厦门大学学报: 自然科学版, 2003, 42: 629-632.

徐磊, 夏宁, 姜学钧. 青岛地区地表天然放射性调查[J]. 核农学报, 2011, 25(4): 773-778.

薛禹群, 谢春红, 吴吉春等. 海水入侵、咸淡水界面运移规律研究[M]. 南京: 南京大学出版社, 1991.

薛禹群, 谢春红, 吴吉春. 海水入侵研究[J]. 水文地质工程地质, 1992, 6: 29-33.

王大纯, 张人权, 史毅虹等. 水文地质学基础[M]. 北京: 地质出版社, 2002.

王芬芬. 应用镭同位素二维平流扩散模型计算黄东海水体的涡动扩散和平流速率[D]. 厦门大学硕士论文, 2011.

王寿景, 李立, 李燕初等. 厦门浔江湾海洋水文特征[J]. 台湾海峡, 1998, 17(2): 201-206.

王绍鸿, 杨建明, 俞鸣同等. 福建沿海湾更新世以来海平面变化[J]. 台湾海峡, 1994, 13(2): 166-175.

吴彩容, 刘剑秋, 王绍鸿等. 福建漳浦前湖湾硅藻组合及其古环境意义[J]. 海洋通报, 2008a, 27(1): 75-80.

吴彩容, 刘剑秋, 王绍鸿等. 福建漳浦前湖湾沉积物磁性性质特征分析[J]. 华侨大学学报(自然科学版), 2008b, 29(1): 111-115.

吴吉春, 薛禹群, 谢春红等. 海水入侵过程中水 - 岩间的阳离子交换[J]. 水文地质工程地质, 1996, 3: 18-19.

谢永臻. 南海及厦门邻近海域镭同位素地球化学的研究 [D]. 厦门大学博士学位论文, 1994, 10-18.

谢永臻, 黄奕普, 施文远等. 天然水体中 ^{226}Ra 、 ^{228}Ra 的联合富集与测定[J]. 厦门大学学报, 1994, 33: 86-90.

姚庆元, 何昭星, 庄进耀. 厦门岛晚第四纪沉积层的划分与海平面变化的关系[J]. 华南地震, 1988, 8(4): 25-33.

姚庆元, 何昭星. 厦门岛地貌类型的划分及其开发利用[J]. 台湾海峡, 1988, 7(4): 395-401.

姚颂恩. 福州市区地貌, 水文特点与防洪[J]. 福建师范大学学报(自然科学版), 1994, 10(4): 104-110.

于静洁, 宋献方, 刘相超等. 基于 D 和 ^{18}O 及水化学的永定河流域地下水循环特征解析[J]. 自然资源学报, 2007, 22(3): 415-423.

袁晓婕. 镭同位素示踪胶州湾的海底地下水排泄[D]. 厦门大学博士学位论文, 2013.

章斌, 郭占荣, 高爱国等. 用氢氧稳定同位素评价闽江河口区的地下水输入[J]. 水科学进展, 2012, 23(4): 539-548.

章斌, 郭占荣, 高爱国等. 用氢氧稳定同位素揭示闽江河口区河水, 地下水和海水的相互作用[J]. 地球学报, 2013, 34 (2) : 213-222.

章光新, 何岩, 邓伟. 同位素 D 与 ^{18}O 在水环境中的应用研究进展[J]. 干旱区研究, 2004, 21(3): 225-229.

张文开. 福州城市地貌与城市气候关系分析[J]. 福建师范大学学报(自然科学版), 1998, 14(4): 96-102.

章新平, 刘晶淼, 中尾正义等. 我国西南地区降水中过量氙指示水汽来源[J]. 冰川冻土, 2009, 31(4): 613-619.

章新平, 姚檀栋. 我国降水中 ^{18}O 的分布特点[J]. 地理学报, 1998, 53(4): 356-364.

张章新. 闽江流域水文特性分析[J]. 水文, 2000, 20(6): 55-58.

郑志强. 福州地铁规划建设中的水文地质环境问题分析[J]. 绿色科技, 2013, 11: 153-157.

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库